

Anti-Wrinkle Treatment

Advice is given to ensure the client fully understands what should and should not be done, carried out or will be experienced post treatment to achieve maximum treatment results and to prevent risk of infection.



Pre treatment advice:

- Do NOT consume alcoholic beverages at least 24 hours prior to treatment (alcohol may thin the blood and increase the risk of bruising)
- Avoid anti-inflammatory/blood thinning medications, if possible for a period of 2 weeks before treatment. Medications and supplements such as aspirin, vitamin E, ginkgo biloba, ginseng, St. John's Wort, Omega 3/Fish Oil supplements, Ibuprofen, Motrin, Advil, Aleve and other NSAIDS have a blood thinning effect and can increase the risk of bruising and swelling after injections.
- Schedule your Dermal Filler and Botox appointment at least 2 weeks prior to a special event which you may be attending, such as a wedding or a vacation. Results from the Dermal Filler and Botox injections will take approximately 4 to 7 days to appear. Also bruising and swelling may be apparent in that time period. Sculptra does take longer to see results so plan accordingly with your provider.
- Discontinue Retin-A 2 days before and 2 days after treatment.
- Reschedule your appointment at least 24 hours in advance if you have a rash, cold sore or blemish on the area.
- If you have a history of cold sores please let your provider know, they may put you on an anti-viral medication prior to treatment.
- Be sure to have a good breakfast, including food and drink before your procedure. This will decrease the chances of lightheadedness during your treatment.
- You are not a candidate if you are pregnant or breastfeeding.

Aftercare advice:

- Do NOT manipulate the treated area for 3 hours following treatment. Do NOT receive facial/ laser treatments or microdermabrasion after Botox injections for at least 10 days. Ask your provider if you are not sure about the time frame of certain services.
- Some providers believe that smiling and frowning right after Botox treatments helps the Botox find its way to the muscle into which it was injected after treated.
- Do NOT lie down for 4 hours after your Botox treatment. This will prevent the Botox from tracking into the orbit of your eye and causing drooping eyelid.
- It can take approximately 4 to 7 days for results to be seen. If the desired result is not seen after 2 weeks of your treatment you may need additional Botox. You are charged

for the amount of product used. Therefore, you will be charged for a product used during any touch-up or subsequent appointments.

- Do NOT perform activities involving straining, heavy lifting, or vigorous exercise for 6 hours after treatment. This will keep the Botox in the injected area and not elsewhere.

Please make sure that you fully understand everything, if you are unsure about anything or have any further questions please feel free to contact me by phone or email.

Suzanne Seth

Tel: 01383 871744

Email: suzanne@hiddenbeauty.info